

AND A CATEGORIFICATION OF THE FREED-QUINN LINE BUNDLE

joint work with Dan Berwick-Evans

Laura Murray

Apurva Nakade

Emma Phillips

Thanks to the ANS NRC on geometric representation theory
and equivariant elliptic cohomology.

Plan: 1) Context / Motivation

2) Principal 2-group bundles: definitions and key properties

3) Applications

§ 1. CONTEXT AND MOTIVATION

Let G be a compact Lie group and fix $\alpha \in H^3(G, U(1))$

→ there are lots of fun things we can do with this data.

- • Chern-Simons theory
- G -equivariant elliptic cohomology
- • String structures
- Equivariant gerbes
- Higher representation theory
- ⋮

★ Smooth two-group $\mathfrak{G} = \mathfrak{G}(G, U(1), \alpha)$

and its moduli space of principal 2-group bundles

Expectation

All of the above topics are meaningfully related to the space $\mathrm{Bun}_{\mathfrak{G}}(X)$.

- Today we focus on the case that G is finite.

Fix input data: G - finite group

A - an abelian Lie group with trivial G -action
(e.g. $A = U(1)$)

$\alpha: G^{*3} \rightarrow A$ a 3-cocycle.

Chern-Simons story: ($A = U(1)$)

↳ Dijkgraaf-Witten theory.

Freed-Quinn: construct a line bundle \mathcal{L} on the moduli space of principal G -bundles on Riemann surfaces.

- For X a Riemann surface, we get a line bundle \mathcal{L}_X on $Bun_G(X)$.
- $\Gamma(Bun_G(X), \mathcal{L}_X)$ is exactly the vector space that Chern-Simons assigns to X .

Elliptic aside Restricting \mathcal{L} to the moduli space of principal G -bundles over elliptic curves, we obtain $\tilde{\mathcal{L}}$.

Ganter: $\tilde{\mathcal{L}}$ is the natural home of twisted G -equivariant elliptic cohomology.

2-groups story $[\alpha] \in H^3(G; A)$ classifies finite 2-groups

that are central extensions of G by $*//A$.

- To the representative α , we associate \mathcal{G} with
 - objects $g \in G$
 - morphisms $\text{Hom}_{\mathcal{G}}(g, h) = \begin{cases} \emptyset & g \neq h \\ A & g = h \end{cases}$
- \otimes -structure: $g \otimes h = gh$.

$$\text{associativity: } (g \otimes h) \otimes k \xrightarrow{\sim} g \otimes (h \otimes k)$$

$\uparrow \alpha(g, h, k) \in A$

- \mathcal{G} can be viewed as a smooth 2-group, depending on the smooth structure of A .

(i.e. a group object in Bibun)

- Today
- we'll define a moduli space (bicategory) $\text{Bun}_{\mathcal{G}}(X)$
 - we'll see there is a nice map $\text{Bun}_{\mathcal{G}}(X) \rightarrow \text{Bun}_G(X)$
 - when X is a Riemann surface, this categorifies \mathcal{L}_X .
 - Sections of \mathcal{L}_X are then isomorphism classes of lifts from $\text{Bun}_G(X)$ to $\text{Bun}_{\mathcal{G}}(X)$.
 - also applications to string structures/string geometry.

§2. Principal 2-group bundles

Fix a smooth (finite) 2-group \mathcal{G} and smooth manifold.

Goal Define a bicategory of smooth 2-group bundles over X

Our favourite definition: A principal \mathcal{G} -bundle on X is a smooth stack $\mathcal{P} \rightarrow X$ equipped with an action of \mathcal{G} which is locally trivial:

\exists surjective submersion $u: Y \rightarrow X$ and an isomorphism

of \mathcal{G} -stacks over Y :

$$d: u^* \mathcal{P} \xrightarrow{\sim} Y \times_{\mathcal{G}} \mathcal{G}.$$

Cech data for \mathcal{P} - tells us how to glue \mathcal{G} -bundles from the trivial bundle on an open cover \mathcal{Y} .

Note: the first level of gluing data is a bibundle Φ

$$Y \times Y \times \mathcal{G} \xrightarrow{\Phi} Y \times Y \times \mathcal{G}$$

which we can assume is trivial as an A -bundle)

- $u: Y \rightarrow X$
- $g: \underset{x}{Y \times Y} \rightarrow G$ satisfying "cocycle" conditions.
- $\gamma: \underset{x}{Y \times Y \times Y} \rightarrow A$

Example: An A -gerbe over X is a principal $*//A$ -bundle over X .

Cech data: $u: Y \rightarrow X$
 $\gamma: \underset{x}{Y \times Y \times Y} \rightarrow A$ a 2-cocycle.

Definition An A -2-gerbe is determined by

$$u: Y \rightarrow X$$

$$\lambda: \underset{x}{Y \times Y \times Y \times Y} \rightarrow A \text{ a 3-cocycle.}$$

Observe We have a forgetful functor $\pi: \text{Bun}_G(X) \rightarrow \text{Bun}_G(X)$
in terms of Cech data: $(u, g, \gamma) \mapsto (u, g)$

Theorem [Berwick-Evans, C, Murray, Nakade, Phillips]

$\pi: \text{Bun}_G(X) \rightarrow \text{Bun}_G(X)$ is a tensor over the symmetric monoidal
bicategory $\text{Gerbe}_A(X)$.

Sketch of proof The fibre over $\overset{P''}{(u, g)}$ is given by the γ 's that
complete the triple (u, g, γ) .

- P determines a 3-cocycle $\lambda_{P,\alpha} = g^* d: Y_x^4 \rightarrow A$:

$$(y_1, y_2, y_3, y_4) \mapsto \alpha(g(y_1, y_2), g(y_2, y_3), g(y_3, y_4))$$


i.e. a 2-gerbe.

Claims 1) the data of γ is equivalent to a trivialisation
of this 2-gerbe $\lambda_{P,\alpha}$

2) the bicategory of trivialisations of a fixed 2-gerbe
is a torsor over the symmetric monoidal bicategory
of gerbes. \square

Principal 2-group bundles in terms of classifying stacks:

Expectation: \mathbb{G} -bundles are classified by maps $X \rightarrow B\mathbb{G}$.



Definition: A flat \mathbb{G} -bundle is a principal \mathbb{G}_{dis} -bundle
 \hookrightarrow discrete topology.

Recall: Flat principal G -bundles are classified by homomorphisms

$$\pi_1(X) \rightarrow G$$

Theorem [BCMNP]

For X with contractible universal cover:

$$\begin{array}{ccc} \text{Bun}_{\mathbb{G}}^b(X) & \xrightarrow{\sim} & \text{Hom}_{\text{Bicat}}(*//\pi_1(X), *//\mathbb{G}) \\ \downarrow \pi & & \downarrow \pi \\ \text{Bun}_G^b(X) & \xrightarrow{\sim} & \text{Hom}_{\text{Cat}}(*//\pi_1(X), *//G) \cong \text{Hom}_{\text{Grp}}(\pi_1(X), G) // G \end{array}$$

A homomorphism $\pi_1(X) \rightarrow \mathbb{G}$:

- $g: \pi_1(X) \rightarrow G$ a homomorphism
- For all $a, b \in \pi_1(X)$. $g(a)g(b) \xrightarrow{\sim} g(ab) \in \mathbb{G}$
 $\gamma(a, b) \in A$ + cocycle condition

A natural transformation $(g_1, \gamma_1) \Rightarrow (g_2, \gamma_2)$:

- $t \in G$ s.t. $\forall a \in G, t g_1(a) t^{-1} = g_2(a)$
- $\forall a \in G, t g_1(a) \xrightarrow{\sim} g_2(a)t \in \mathcal{G}$
 \uparrow
 $w(a) \in A$ + cocycle condition
- 2-morphisms are given by $w \in A$.

Theorem [BCMNP] The action of G on $\text{Hom}_{\text{grp}}(\pi_1(X), G)$ lifts to an action of G on the bicategory $\text{Hom}_{\text{bicat}}(*//\pi_1(X), *//\mathcal{G})$.
This gives TC the structure of a cloven 2-fibration.

§3. APPLICATIONS

3.A. Freed-Quinn line bundle ($A = U(1)$, X a Riemann surface)

- We've seen that $\text{TC}: \text{Bun}_G^b(X) \rightarrow \text{Bun}_G^b(X)$ is a cloven 2-fibration with fibres equivalent to $\text{Gerbe}_{U(1)}(X)$
- We can now take isomorphism classes along fibres.
 - Since isomorphism classes of gerbes are given by $H^2(X, U(1)) \cong U(1)$,

we obtain a principal $U(1)$ -bundle on $\text{Bun}_G^b(X)$.

Theorem [BCMNP] The associated line bundle is the Freed-Quinn line bundle \mathcal{L}_X .

(cf. Willerton for X a torus)

§4.B string structures

- We work with the string 2-group (not finite!)

$$1 \rightarrow *//U(1) \longrightarrow \text{String}(n) \longrightarrow \text{Spin}(n) \longrightarrow 1.$$

- Starting from a finite group representation $g_0: G_0 \rightarrow SO(n)$, we produce a 2-group:

$$\begin{array}{ccccc}
 *//_{U(1)} & \longrightarrow & \mathbb{G} & \longrightarrow & \text{String}(n) \\
 \downarrow & & \downarrow & & \downarrow \\
 \mathbb{Z}/2\mathbb{Z} & \longrightarrow & G & \xrightarrow{g} & \text{Spin}(n) \\
 \downarrow & & \downarrow & & \downarrow \\
 G_0 & \xrightarrow{g_0} & & & SO(n)
 \end{array}$$

Recall Let $P_0 \rightarrow X$ be an oriented vector bundle with structure group G .

↳ a spin structure on P_0 is a lift to a G -bundle P .

Definition [Schommer-Pries] a string structure on P is a lift to a \mathbb{G} -bundle \mathcal{P}

Alternative definition [Waldorf] a string structure on P is a trivialization of the 2-gerbe $\lambda_{P,\alpha}$

Theorem [BCMNP] The two definitions coincide.

- Let $P \rightarrow X$ be a principal flat G -bundle and consider Chern-Simons theory for P , C_S .

Definition [Stolz-Teichner] a geometric string structure on P is a trivialization of C_S .

- in particular, for suitable $f: M^2 \rightarrow X$, $C_S(f)$ is a line, and we require a non-zero point in this line.

i.e. an isomorphism class of flat \mathbb{G} -bundle over $f^*\mathcal{P}$

- this could be given by $f^*\mathcal{P}$ for \mathcal{P} a flat lift of P .

\mathcal{S} (part of the data of a trivialization of \mathcal{CS}_P)



(part of the data of $\mathcal{D} \in \pi^{-1}(P) \subset \text{Bun}_{\mathcal{G}}^b(X)$)

Work in progress: complete this story.

Thank you!